

**ASPIRIN- aspirin tablet, film coated**  
**Topco Associates, LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**TopCare 44-157**

***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

temporarily relieves:

- minor pain of arthritis
- headache
- menstrual pain
- muscle pain
- toothache
- pain and fever of colds

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems

- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- you are taking a diuretic
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- drink a full glass of water with each dose

- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

**Inactive ingredients**

corn starch, hypromellose, polyethylene glycol, propylene glycol

**Questions or comments?**

**1-888-423-0139**

**Principal display panel**

TopCare®  
health™

NDC 36800-157-12

COMPARE TO GENUINE BAYER® ASPIRIN  
ACTIVE INGREDIENT†

**REGULAR STRENGTH**

**Aspirin 325 mg**

PAIN RELIEVER • FEVER REDUCER

**(NSAID)**

- Safe Pain Relief • Easy to Swallow
- Sodium & Caffeine Free

**100 COATED TABLETS**

• OUR PHARMACISTS RECOMMEND •

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

DISTRIBUTED BY TOPCO ASSOCIATES LLC, ELK GROVE VILLAGE, IL 60007

©TOPCO LNKA0818 QUESTIONS? 1-888-423-0139

topcare@topco.com www.topcarebrand.com

**Visit here for more information:**

**<http://topbrnds.com/48ZS2N>**

†This product is not manufactured or distributed by  
Bayer HealthCare LLC, owner of the registered trademark  
Genuine Bayer® Aspirin.

50844 REV0318A15712

Adhesive Only

STOP PEELING

### Drug Facts (continued)

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer  
■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if** ■ you are taking a diuretic  
■ stomach bleeding warning applies to you

■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
■ you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for** ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**

■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood  
■ have bloody or black stools

■ have stomach pain that does not get better  
■ ringing in the ears or a loss of hearing occurs

■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days

■ redness or swelling is present ■ new symptoms occur  
These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy, unless definitely directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions** ■ drink a full glass of water with each dose  
■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor  
■ children under 12 years: do not use unless directed by a doctor

### Other information

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

**Inactive ingredients** corn starch, hypromellose, polyethylene glycol, propylene glycol

**Questions or comments?** 1-888-423-0139

**TopCare**  
health™

NDC 36800-157-12

COMPARE TO GENUINE BAYER® ASPIRIN  
ACTIVE INGREDIENT\*

REGULAR STRENGTH

**Aspirin**  
**325 mg**

PAIN RELIEVER • FEVER REDUCER  
(NSAID)

- Safe Pain Relief • Easy to Swallow
- Sodium & Caffeine Free

**100** COATED TABLETS



Visit here for more information:  
<http://topcarebrands.com/48252N>

4 21701 0089 0

This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Genuine Bayer® Aspirin. 50844 REV0318A15712  
DISTRIBUTED BY TOPCO ASSOCIATES LLC, ELK GROVE VILLAGE, IL 60007  
©TOPCO LINKA0818 QUESTIONS? 1-888-423-0139  
topcare@topco.com www.topcarebrand.com

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

### Drug Facts

#### Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)\* ..... Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

#### Uses temporarily relieves:

- headache ■ muscle pain ■ minor pain of arthritis
- toothache ■ pain and fever of colds ■ menstrual pain

#### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ facial swelling ■ shock ■ hives ■ asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older

■ take a blood thinning (anticoagulant) or steroid drug  
■ have 3 or more alcoholic drinks every day while using this product ■ have had stomach ulcers or bleeding problems

■ take more or for a longer time than directed  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

PEEL HERE FOR MORE DRUG FACTS

44-157

## ASPIRIN

aspirin tablet, film coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-157
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	

<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;157;ASPIRIN
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-157-87	1 in 1 CARTON	02/10/1996	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:36800-157-12	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/1996	
3	NDC:36800-157-14	500 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/1996	
4	NDC:36800-157-17	300 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/1996	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	02/10/1996	

**Labeler** - Topco Associates, LLC (006935977)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(36800-157)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(36800-157)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	pack(36800-157)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(36800-157)

Revised: 4/2021

Topco Associates, LLC